
(Pinus nigra)

(Picea excelsa)

)
/ m / Cm)

×
(
(/ m / Cm

...
.()

. / m / Cm

.()

) ()
m / Cm (/

.()

.()

.() (/ m)

/ Cm

.() / m

/ m / Cm

.()

	p.p.m	p.p.m		O.C	PH	ECe	S.P	Cm	
		/	/	/		/	/		
		/	/	/	/	/			
		/	/	/	/	/	/		
		/	/	/	/	/	/		
		/	/	/	/	/	/		

(Pinus nigra)

Corylus)

(Picea excelsa)

(avellana

...

		(m)		(ha)		

∴
() T-test ×
() ()
×

Levenes

T-test (×)
/
/
∴
∴
∴

±

p	t	df	± Sd	
/ *	/			
			/ ± /	/ ± /

/

/

±

p	t	df	± Sd	
/ **	/			
			/ ± /	/ ± /

(/)

/

±

p	t	df	± Sd	
/ Ns	/			
			/ ± /	/ ± /

p = / *
p = / **
p = / ns

...

()

/ m / Cm)
(/ m / Cm

.()

()

/ Cm)
(/ m

)

(

.()

()

)
(/ Cm)
(/ m

(*Picea abies* (L.) Karst)

Picea excelsa

6-Daniel B. Houston., James R. 1996. Evaluation of Sphaeropsis Blight Impact on Geographic Seed Sources of *Pinus nigra* in Ohio. Ohio Department of Natural Resources Division of Forestry. 4 p.

7- Koprivica- M ., Ratknic-M ., Markovic- N. 1998. Growth and development trees in artificially established coniferous stands in Ivanjica region (Serbia, Yugoslavia). Zbornik – radova – Institut – za – Sumarstvo (Yugoslavia) . v. 42-43 p . 47-58.

8- Veselinovica – N., Markovic – D., Peno – M., Vuletic – D. 1984. Analysis of the development of black pine plantation on sandy soils, disturbed by the surface coal mining in REIK Kolubara (Yugoslavia). Zemljiste – i – biljka (Yugoslavia) . V. 35(1) p . 67-74.

A Quantitive and Qualitive Study of *Picea excelsa* and *Pinus nigra* in Fandoghloo Forest Reserve.

F. Gh. Aghbash¹ Gh. A. Jalali² A. Teymourzadeh³ M. Hosseini⁴

Abstract

To study the quantitative and qualitative characteristics of *Picea excelsa* and *Pinus nigra* plantations in Ardabil, Fandoghloo forest reserve was selected. Seventeen square sampling plots (10×10 m) were selected using randomized systematic sampling. Quantitative characteristics included survival rate, total and height, collar diameter while vitality was taken as the qualitative characteristic. A comparative study between *Picea excelsa* and *Pinus nigra* on identical aspect and slope (southern , 20 %) indicated a higher growth of diameter and height for *Picea excelsa*. (*Picea excelsa* 1.12 Cm and 0.26 m , *Pinus nigra* 0.64 Cm and 0.15 m). Results showed that from survival point view, there was no significant difference between the species. From vitality point of view, *Picea excelsa* was superior.

Keywords: Plantation, Quantitive and Qualitive Study, *Picea excelsa*, *Pinus nigra*, Fandoghloo, Ardabil.

1- M. Sc, Tarbiat Modarres University, E-mail: ghasemifarhad@yahoo.com

2-Assistant Professor, Tarbiat Modarres University

3- Assistant Professor, Mohaghegh e Ardabili University

4- Assistant Professor, Tarbiat Modarres University