



Original research

Anti-carcinogenic effects of *Satureja khuzistanica* and *Zataria multiflora* essential oils on K562 cell line proliferation

Shller Khakzad^a, Fatemeh Rahmani^a, Mohammad Hojjati^{b,*}, Mohammad Reza Tabandeh^c^a Department of Biology, Faculty of Sciences, Urmia University, Urmia, Iran^b Department of Food Science and Technology, Agricultural Sciences and Natural Resources University of Khuzestan, Ahvaz, Iran^c Department of Biochemistry and Molecular Biology, Faculty of Veterinary Sciences, Shahid Chamran University of Ahvaz, Iran

ABSTRACT

The aim of this study was to investigate cytotoxic effect of *Satureja khuzistanica* and *Zataria multiflora* essential oils on PBMC and K562 cell lines growth inhibition. The plant essential oils were extracted by hydro-distillation and analyzed by GC-MS. In addition, the total phenol content and antioxidant activity of oils were assessed. The PBMC and K562 cell lines were cultured in RPMI medium containing 10% FBS supplemented with 0, 12.5, 25, 50 and 100 (µg/ml) of essential oils for 24, 48 and 72 h. The MTT assay revealed anti-proliferative effect of essential oils on both cell lines in a dose dependent manner. These effects became more prominent after 72 h. Both essential oils contained high amounts of phenolic compounds such as thymol and carvacrol. Due to the higher percentage of carvacrol, which has more toxicity than thymol, *S. khuzistanica* had a greater antioxidant activity as well as inhibitory effect on cell growth. In conclusion, the *S. khuzistanica* and *Z. multiflora* essential oils have the potential to be used in the production of anticancer medicines.

Keywords: Anti-proliferation, Antioxidant activity, Carvacrol, Cytotoxicity, GC-MS

Received 19 April 2019; Received 7 July 2019; Accepted 16 July 2019

1. Introduction

Cancer is an uncontrolled cell proliferation leading to second cause of mortality worldwide (Hassanpour & Dehghani, 2017). The chemicals, radiations, unhealthy diet, environmental factors, infection and tobacco smoke are risk factors for cancer (Anand et al., 2008). The environmental carcinogenic substances influence directly or indirectly the cell cytoplasm and nucleus (Poon et al., 2016). Cancer occurs when mutations in vital genes lead to dysfunction of genes, disruption of cell cycle and abnormal proliferation (Aizawa et al., 2016). Overall, the cancer prevalence worldwide has raised over the past decade; for instance; 1,665,540 and 585,720 Americans were diagnosed and died from cancer in 2014 (Hassanpour & Dehghani, 2017).

Traditional herbal medicine has become center of attention during the last decades and number of member state regulating herbal medicine has increased from 65 (1999) to 119 (2012). Moreover, the research institutes for herbal medicine has been upgraded from 19 (1999) to 73 (2012) (W.H.O, 2013). There are hundreds of literature data revealing the importance and utilization of herbal medicines to cure diseases like cancer (Rafieian-Kopaei et al., 2013). According to latest WHO reports, traditional system of herbal treatment has even been adapted in developed countries

including Belgium (31%), Australia (48%), France (49%), Canada (70%) and Germany (77%) (WHO, 2013).

Natural products are pharmacologic agents and have been traditionally used for prevention and cure of chronic diseases (Sefidkon et al., 2013). Today, blood cancer is one of the most important diseases and plants are potential candidates for cancer treatment (Zali et al., 2011). Aromatic plants contain essential oils with chemo-preventive properties due to volatile extracts comprising of a wide variety of active phytochemicals, such as monoterpenes, polyphenols and flavonoids (Lai & Roy, 2004).

Satureja khuzistanica Jamzad (Persian name, 'marzeh khuzistani') belongs to the family Lamiaceae, and is an endemic plant of Iran with wide distribution in the southern part of this country. The plant has commercial and medicinal values and has been widely used in pharmaceutical, food, cosmetics and perfumery industries due to its high essential oil, which mainly contain carvacrol and terpinene compounds (Kouravand et al., 2018; Sefidkon et al., 2013). In traditional medicine, *S. khuzistanica* has been used as an antimicrobial, spasmolytic cicatrisant, diuretic (Gohari et al, 2005), analgesic, antiseptic and anti-inflammatory agent (Naghbi et al., 2005). Ghazanfari et al. (2006) demonstrated that oral consumption of *S. khuzistanica* essential oil has remarkable impact in the treatment of inflammation and diabetes.

*Corresponding author.

E-mail address: hojjatim@yahoo.com (M. Hojjati).

Zataria multiflora Boiss (vernacular name of Avishan Shirazi, in Iran) belongs to the family Lamiaceae and geographically grows in Iran, Pakistan and Afghanistan (Hosseinzadeh et al., 2000). The plant has traditionally been used as an antiseptic, anesthetic and anti-spasmodic drug (Ali et al., 2000), also extensively used as a flavor ingredient in Iranian foods. The flowered browses of the plant contain essential oil with various pharmaceutical, antimicrobial, and antioxidant effects (Saeedi-Saravi et al., 2008). The main constituents of its essential oil are phenolic compounds such as carvacrol and thymol (Zandi-Sohani & Ramezani, 2015). The cytotoxicity and antibacterial capacity (Amin et al., 2010) as well as radioprotectivity and chemoprotectivity (Hosseini-mehr et al., 2011) effects of *Z. multiflora* essential oils have been reported.

To the Knowledge of authors, no research has been conducted on the cytotoxic effects of *S. khuzistanica* and *Z. multiflora* plant extracts on cell viability of K562 leukemia cell line. Therefore, we aimed to investigate the apoptotic effects of these two species extract on cell viability of PBMC and K562 cells.

2. Material and Methods

2.1. Extraction and analysis of essential oil

The aerial parts of *S. khuzistanica* were collected from Khuzestan province in the southwest of Iran during full flowering stage in 2017. The aerial parts of *Z. multiflora* were harvested in pre-flowering stage from wild grown plants in the Mountain regions of Fars province located in the south of Iran. The harvested plants were dried at room temperature for two weeks, then, air-dried plants (in each habitat) were ground and powdered with mixer for essential oil extraction and other experiments. The powdered plants aerial parts (250 g) were hydro-distilled using a Clevenger type apparatus for 3 h, according to the method recommended in the European Pharmacopoeia (Council of Europe, 1997). The resultant essential oil was dried over anhydrous sodium sulfate and stored at 4°C until analyzed and tested. Dilutions of the oil in the various assays were then made from the parent oil stock by addition of dimethyl sulfoxide (0.1%; DMSO) to yield the desired final test concentrations (Hojjati & Barzegar, 2017).

2.2. GC-MS analysis

The chemical analysis of the essential oils was performed using Agilent technology GC-6890A and GC-MSD 5975 fitted with a HP-5MS column (30 mm × 0.25 mm i.d, 0.25 mm film thick). 0.5 µL of essential oils were injected for analysis, and Helium was used as carrier gas at 1 mL/min. The oven temperature was programmed from 60°C to 250°C at the 5°C/min rate, and was finally hold for 1 min. Injector and detector temperatures were 230°C and 300°C, respectively. The compounds of essential oils were identified according to GC-MS Retention time, Kovats index and Mass spectra NIST, 2013 (National Institute of Standards and Technology, 2013).

2.3. Total phenolic content and antioxidant activity of essential oils

Total phenolic content of the essential oils was measured according to the absorbance values of the oil reacted with Folin-

Ciocalteu reagent at 765 nm and compared the results with Gallic acid as standard solution according to the method described previously with minor modifications (Hojjati & Barzegar, 2017). Total phenolic content was expressed as milligrams of Gallic acid equivalent per gram of essential oil (mg GAE/g). Antioxidant activity of the essential oils was assessed using DPPH and ferric reducing antioxidant power (FRAP) assays based on original methods by Hojjati and Barzegar (2017) and Saidi (2014) with some modifications, respectively.

2.4. Cell culture

One type of cancer cells (K562) and one type of normal cells (PBMCs) were used to determine the cell viability against essential oils exposure. The human myeloid leukemia K562 cells were obtained from Pasture Institute of Iran, Tehran, Iran. Fresh peripheral blood was taken with informed consent from three healthy volunteers into sterile falcon tubes containing heparin (200 IU/ml). Human peripheral blood mononuclear cells (PBMC) were isolated by using 1.077 g/ml Ficoll/ Hypaque as previously described (Boyum, 1968).

Culture medium used for the growth of K562 was as follows: RPMI 1640 (Gibco, Life technologies GmbH, Frankfurt, Germany) enriched with Fetal Bovine Serum-FBS 10%, Streptomycin (100 µg/ml, Sinagen, Tehran, Iran) and Penicillin (100 µg/ml, Sinagen, Tehran, Iran). The cells were grown in incubator (holding British Standard) with 5% CO₂, 95% humidity and 37°C (Aslani et al., 2014).

2.5. Assessment of cytotoxicity based on MTT

Growth of cells was quantitated by the ability of living cells to reduce the yellow dye 3-(4, 5-dimethyl-2-thiazolyl)-2, 5-diphenyl-2H-tetrazolium bromide (MTT) to a blue formazan product. MTT assay is an accurate method to evaluate the survival of cells. This assay is based on the cleavage of the yellow tetrazolium salt, MTT, to form a soluble blue formazan product by mitochondrial enzymes, and the amount of formazan produced is directly proportional to the number of living cells (Sylvester, 2011). In this study, to evaluate the effects of *Z. multiflora* and *S. khuzistanica* essential oils on the morphology of PBMC and K562 cell line, 104 cells were positioned in 96-well plates (Surface, Denmark). The concentrated plant essential oil was dissolved in dimethyl sulphoxide (DMSO) (Sigma Aldrich, Germany) to obtain appropriate solutions of the essence. The substock solutions of 12.5, 25, 50 and 100 µg/ml was prepared by diluting the stock solution into serum-free culture medium, RPMI 1640 (the percentage of DMSO in the experiment should not exceed 1).

After 24 h, the cells were treated with different concentrations of *S. khuzistanica* and *Z. multiflora* essential oils (12.5, 25, 50, 100 µg/ml), prepared separately. The normal cells and K562 cell line were incubated for 24, 48 and 72 h with different concentrations of essential oils (12.5, 25, 50, 100 µg/ml). Then, 20 µl MTT solution 0.5 mg/ml was added into the wells and incubation continued at 37°C for 3 to 5 h. After this time, the supernatant was removed and 200 µl of DMSO solution was added at room temperature for 15 min. Finally, the percentage of cell viability was determined by ELISA reader at 570 nm. The percentage of cell viability was calculated using the following equations:

[mean optical density (OD) of treated cells/mean OD of control cells] × 100. Percent of cytotoxicity = 100 – percent of Viability

(Zamanian-Azodi et al., 2012). The effects of essential oil were expressed by IC50 values (the drug concentration reducing the absorbance of treated cells by 50% with respect to untreated cells).

Table 1. Volatile composition of *S. khuzistanica* and *Z. multiflora* essential oils.

Compound	*RT (min)	*Kl _{exp}	%	
			<i>Satureja</i>	<i>Zataria</i>
α- thujene	5.253	927	2.34	0.65
α- pinene	5.420	934	2.67	3.32
Camphene	5.742	947	0.27	0.25
β- pinene	6.375	972	1.81	0.66
beta myrcene	6.664	984	2.44	1.74
3- octanol	6.80	993	-	0.19
α- phellandrene	7.009	997	0.70	0.24
delta 3 carne	7.153	1005	0.16	0.07
α- terpinene	7.331	1011	4.56	2.28
p- cymene	7.575	1019	9.55	7.34
limonene	7.642	1022	1.42	0.83
1,8 cineole	7.775	1026	0.29	0.68
β-ocimene	8.097	1037	0.20	0.07
γ- terpinene	8.486	1050	22.72	6.66
(z)-sabinene hydrate	8.742	1058	0.13	0.21
cis linalool oxide	8.845	1072	-	0.10
α- terpinolene	9.175	1081	0.19	0.25
trans linalool oxide	9.267	1088	-	0.08
linalool	9.697	1090	0.06	5.26
hotrienol	9.825	1098	-	0.18
borneol	11.830	1168	0.57	0.25
terpinene-4-ol	11.941	1176	0.59	1.13
α- terpineol	12.108	1184	0.06	1.01
cis dihydrocarvone	12.555	1200	-	0.07
thymyl methyl ether	13.111	1224	-	1.27
carvacrol	13.352	1227	0.21	1.88
methylether				
carvone	13.852	1243	1.12	-
anethole	14.497	1258	0.39	-
thymol	14.674	1279	1.82	30.61
carvacrol	15.563	1300	38.33	22.18
piperitenone oxide	16.741	1363	0.11	-
carvacryl acetate	16.830	1370	0.80	1.13
carvacryl acetate	16.844	1380	-	1.13
caryophyllene	18.029	1418	0.79	2.56
β-gurjenene	18.232	1432	-	0.07
alloaromadendrene	18.507	1449	0.21	0.11
aromadendrene	18.521	1449	-	1.17
α- humulene	18.803	1481	0.06	0.17
ledene	19.874	1500	0.13	0.83
β-bisabolene	20.163	1509	1.70	-
α- amorphene	20.318	1516	0.05	-
delta cadinene	20.518	1520	0.05	-
spathulenol	21.873	1570	0.12	1.03
caryphyllene oxide	22.04	1579	0.47	1.06
isospathulenol	23.243	1640	-	0.13
bisabolene epoxide	23.965	1699	-	0.14
total			97.12	99.14

*RT: Retention Time; Kl_{exp}: Kovats ndex in this experiment; %: Relative percentage obtained from peak area.

2.6. Statistical analysis

All analyses were reported as means ± SD (standard deviation). The SPSS Software v. 24 (SPSS Inc., Chicago, IL, USA) and analysis of variance (ANOVA) were used to investigate significant differences between the average values (95% confidence).

Significant differences between the means were calculated using Duncan test at a significant level of $p < 0.05$.

3. Results and Discussion

3.1. Chemical composition of essential oils

The essential oil obtained by hydro-distillation of the aerial parts of *S. khuzestanica* and *Z. multiflora* was subjected to GC-MS to identify composition. Qualitative and quantitative analytical results of *S. khuzestanica* and *Z. multiflora* essential oils were shown in Table 1. According to gas chromatography–mass spectrometry (GC-MS) analysis, 35 and 40 volatile compounds were identified from *S. khuzestanica* and *Z. multiflora* essential oils representing 97.12% and 99.14% of the total oil, respectively. Based on the Table 1, carvacrol (38.33%), γ terpinene (22.72%) and p-cymene (9.55%) were the main compounds in the *S. khuzestanica* oil. Although Saeidi (2014) reported only carvacrol as the main constituent of *S. khuzistanica* essential oil; Alizadeh (2017) found carvacrol, γ-terpinene, β-bisabolene and pcymene to be the major constituents of the oil of *S. khuzistanica*, which was consistent with our findings. As observed in Table 1, thymol (30.61%), carvacrol (22.18%), Linalool (9.57%), γ terpinene (8.44%) and p-cymene (7.34%) identified as the main components in the essential oil of *Z. multiflora*. It was in close agreement with the previous results (Raiesi et al., 2019; Zandi-Sohani & Ramezani, 2015). Comparison of our findings with others suggests that differences in the components of the essential oils may be attributed to various factors such as the genetic, geographic, cultivar, species, ripening stage, cultural practices, extraction methods, soil and climate conditions (Hojjati & Barzegar, 2017; Saeidi, 2014).

Table 2. Antioxidant activity of *Satureja khuzistanica* and *Zataria multiflora* essential oils.

Essential oil	Total Phenolic Content (mg GA eq/g)	DPPH (%)	FRAP (μmol/g)
<i>Zataria</i>	216.27±17.40 ^b	39.22±4.15 ^b	25.74±2.29 ^b
<i>Satureja</i>	249.53±16.59 ^a	47.19±3.26 ^a	30.02±3.11 ^a

*Values represent mean ± Standard Deviation of three replicates and values with different letters in each column were significantly different at the level of $p < 0.05$.

3.2. Total phenol content and antioxidant activity of essential oils

The results of total phenol content and antioxidant activity of essential oils are shown in Table 2. The level of total phenolic compounds in *S. khuzestanica* and *Z. multiflora* essential oils were 249.53 and 216.27 mg GA eq/g, respectively. In addition, the data showed that in both methods, the antioxidant activity of *S. khuzestanica* oil was significantly higher than that *Z. multiflora* essential ($p < 0.05$). The antioxidant activity of *S. khuzestanica* essential oil in the two methods of DPPH and FRAP assays were 47.19% and 30.02 (μmol/g), respectively, which were significantly higher than the level of antioxidant activity of *Z. multiflora*. The results of the total phenolic content and the antioxidant activity of the essential oils were similar to those of some researchers and were in contrast to some others. The amount of phenolic compounds of *Z. multiflora* essential oil in this study was higher

than the values reported by Mojaddar-Langroodi et al. (2019) and Alizadeh and Shaabani (2014), whereas Raeisi et al. (2019), observed more phenolic compounds. Total phenol content and antioxidant potential of *S. khuzistanica* oil studied in this work was higher than the amount reported by Saidi (2014) and Alizadeh (2017).

As seen in Table 2, the essential oil of *S. khuzistanica* has higher phenolic content and was highest antioxidant activity, compare to *Z. multiflora*. Thus, our results showed that, a correlation between the antioxidant activities and the total phenolic contents was revealed. On the other hand, there was a significant increase in antioxidant activity by increase in total phenolic content of samples. Otherwise, ferric reduction and DPPH radical scavenging occurred more powerfully with the increasing concentrations of phenolic compounds in essential oils. The antioxidant activity of medicinal plants is associated with the phenolic content of them. High redox potentials of phenolic compounds provided them to act as reducing agents, hydrogen donors, and singlet oxygen quenchers (Alizadeh, 2017; Raeisi et al., 2018). Probably because of the different types of phenolic compounds in each essential oil, they showed different antioxidant activity.

3.3. The Effect of essential oils on proliferation of PBMC normal cells and K562 cell line

The results of the effect of four different concentrations (12.5, 25, 50 and 100 µg/ml) of *S. khuzistanica* and *Z. multiflora* essential

oils on cell viability of PBMC and K562 cell line at 24, 48 and 72 h incubation were presented in Fig. 1. Evaluation of cell survival was measured using the MTT test. The 50% growth inhibitory concentration (IC50) of 28.21 and 18.06 (µg/mL) was obtained for *S. Khuzistanica* essential oil for the PBMC and K562 cell line, respectively, and it was found 64.97 and 42.82 (µg/mL) on normal PBMC and K562 cell line, respectively for *Z. multiflora* essential oil.

The absorption rate based on the different concentrations of essential oils at different times was significantly lower than that of the control group (did not receive any essential oil), indicating a toxicity effect of essential oil on PBMC and K562 cell line. According to our results, the amount of cell survival was dependent on the essential oil concentration and incubation time. Hence, the concentration of 100 µg/mL of essential oil and 72 h incubation was the most fatal with the lowest living. Incubation of PBMC from 12.5 to 100 (µg/mL) of *S. Khuzistanica* essential oil for 24 h reduced cell viability from 92.2 to 53.7%, respectively, however, at the same concentrations of essential oil after 72 h of incubation, cell viability was decreased from 78.3 to 47.1%, respectively (Fig. 1A and B). Similarly, at maximum concentration (100 µg/mL) and 72 h incubation, *Z. multiflora* essential oil showed the highest activity on PBMC and K562 cells (Fig. 1C and D). The results showed the effect of both essential oils on the growth of tested cells, although PBMC cell line was less susceptible to the cytotoxicity of both essential oils at the same essential oil concentrations and incubation times compared to the K256 cell line.




Fig. 1. Effects of *S. khuzistanica* (A-B) and *Z. multiflora* (C-D) essential oils on viability of PBMC and K562 cell line.

The presence of large amounts of carvacrol in both essential oils may be due to their anticancer effects because the cytotoxic potential of these monoterpenes have already been proven by Horvathova et al. (2007) who reported that carvacrol and thymol significantly reduced the level of DNA damage induced in K562 cells by the strong oxidant H₂O₂. On the other hand, phenolic compounds such as thymol increase in production of reactive oxygen species (ROS) activity, increase in mitochondrial H₂O₂ production and depolarization of mitochondrial membrane potential (Dutta et al., 2011).

The strong antitumor activity of carvacrol has also been revealed by suppression of mouse B16 melanomas (He et al., 1997), lung tumors in rats (Zeytinoglu et al., 1998) and human larynx carcinoma Hep-2 cells (Stammati et al., 1999).

In addition, the results showed that essential oil of *S. Khuzistanica* had more significant effect on inhibition of cancer cell growth at different incubation times compared to *Z. multiflora*. The results in the present work are in agreement with the findings of Mohammadpour et al. (2015) who investigated the cytotoxicity potential of *S. bakhtiariica* and *Z. multiflora* oils on normal HEK cells compared to MDA-MB-231 and SKOV3 as cancerous cell lines. The results of the analysis of essential oils in this study also indicated that the presence of high levels of terpenoids such as carvacrol, γ -terpinene and p-cymene in *S. khuzistanica* oil.

The efficacy of the plant as an anti-cancer medicine may be related to terpenoid compounds which it could inhibit tumor cell proliferation in human breast and prostate (Yang & Dou, 2010), therefore, the reason for the greater effect of essential oil of *S. Khuzistanica* on cancer cells may be due to the higher amount of phenolic compounds as well as its antioxidant properties compared to *Z. multiflora* (Table 2).

4. Conclusion

Pharmaceutical plants with cytotoxic effect are good candidates to synthesis anticancer drugs. Moreover, due to lower side effects and pharmaceutical intervention, they attract pharmacists' attention to synthesis new drugs in order to treat refractory disease such as cancers (Mongelli et al., 2000). The data presented in this work provided scientific information that might be used for isolation of potentially active compounds from *S. khuzistanica* and *Z. multiflora* medicinal plants in future. Based on our findings, both essential oils had high amount of terpenoids and cytotoxic effect on normal PBMC and K562 cell line due to the presence of high amounts of phenolic compounds such as thymol and carvacrol. However, due to the presence of higher amount of phenolic compounds and antioxidant activity, *S. khuzistanica* pronounced a greater inhibitory effect on cell growth. It is concluded that, the *S. khuzistanica* and *Z. multiflora* essential oils have the potential to be used in the production of anticancer drugs.

Acknowledgment

The authors would like to thank the financial support of the University of Urmia, Iran.

References

Aizawa, K., Liu, C., Tang, S., Veeramachaneni, S., Hu, K. Q., Smith, D. E., & Wang, X. D. (2016). Tobacco carcinogen induces both lung cancer and non-alcoholic steatohepatitis and hepatocellular

carcinomas in ferrets which can be attenuated by lycopene supplementation. *International journal of cancer*, 139(5), 1171-1181.

- Ali, M.S., Saleem, M., Ali, Z., & Ahmad, V.U. (2000). Chemistry of *Zataria multiflora* (Lamiaceae). *Phytochemistry*, 55, 933-936.
- Alizadeh, A. (2017). Essential oil constituents, phenolic content and antioxidant activity of two endemic *Satureja* species from Iran. *Bangladesh Journal of Botany*, 46(3), 925-931.
- Alizadeh, A., & Shaabani, M. (2014). Essential oil composition, total phenolic content and antioxidant activities of Iranian *Zataria multiflora* Boiss. *International Journal of Biosciences*, 4(4), 97-104.
- Amin, M., Kalantar, E., Mohammad-Saeid, N., & Ahsan, B. (2010). Antibacterial effect and physicochemical properties of essential oil of *Zataria multiflora* Boiss. *Asian Pacific Journal of Tropical Medicine*, 3(6), 439-442.
- Anand, P., Sundaram, C., Jhurani, S., Kunnumakkara, AB., & Aggarwal, BB. (2008). Curcumin and cancer: an "old-age" disease with an "age old" solution. *Journal of Cancer Letters*, 267,133-164.
- Aslani, E., Naghsh, N., & Ranjbar, M. (2014). Cytotoxic effects of hydro-alcoholic extracts of cress (*Lepidium Sativum*) – made from different stages of the plant - on k562 Leukemia cell line. *Hormozgan Medical Journal*, 18(5), 370-378.
- Boyum, A. (1968). Separation of leukocytes from blood and bone marrow. Introduction. *Scandinavian Journal of Clinical Laboratory Investigation*, 97, 7-11.
- Council of Europe. (1997). European Pharmacopoeia, 3rd Edition. Strasbourg, France: Council of Europe.
- Dutta, D. D., Parimala, G., Devi, S.S., & Chakraborty, T. (2011). Effect of thymol on peripheral blood mononuclear cell PBMC and acute promyelocytic cancer cell line HL-60. *Chemico Biological Interaction*, 193, 97-106.
- Ghazanfari, G., Minaie, B., Yasa, N., Nakhai, L. A., Mohammadirad, A., Nikfar, S., ... & Salehnia, A. (2006). Biochemical and histopathological evidences for beneficial effects of *Satureja Khuzestanica* Jamzad essential oil on the mouse model of inflammatory bowel diseases. *Toxicology mechanisms and methods*, 16(7), 365-372.
- Gohari, A.R., Hadjiakhoondi, A., Sadat-Ebrahimi, S.E., Saeidnia, S., & Shafiee, A. (2005). Cytotoxic terpenoids from *Satureja macrantha* CA Mey. *Daru- Journal of Faculty of Pharmacy*, 13(4), 177-18.
- Hassanpour, S.H. & Dehghani, M. (2017). Review of cancer from perspective of molecular. *Journal of Cancer Research and Practice*, 4(4), 127-129.
- He, L., Mo, H., Hadisusilo, S., Qureshi, A.A. & Elson, C.E. (1997). Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo. *Journal of Nutrition*, 127, 668- 674.
- Hojjati, M., & Barzegar, H. (2017). Chemical composition and biological activities of lemon (*Citrus limon*) leaf essential oil. *Nutrition and Food Sciences Research*, 4(4), 15-24.
- Horvathova, E., Turcaniova, V., & Slamenova, D. (2007). Comparative study of DNA-damaging and DNA protective effects of selected components of essential plant oils in human leukemic cells K562. *Neoplasma*, 54(6), 478-483.
- Hosseinimehr, S.J., Mahmoudzadeh, A., Ahmadi, A., Ashraf, SA., Shafaghathi, N., & Hedayati, N. (2011). Te radioprotective effect of *Zataria multiflora* against genotoxicity induced by γ irradiation in human blood lymphocytes. *Cancer Biotherapy Radiopharm*, 26 (3), 325-329.
- Hosseinizadeh, H., & Madanifard, M. (2000). Anticonvulsant effects of *Coriandrum sativum* L seeds extracts in mice. *Achieves of Iranian Medicine*, 3,182-184.
- Kouravand, F., Jooyandeh, H., Barzegar, H., & Hojjati, M. (2018). Characterization of cross-linked whey protein isolate-based films containing *Satureja Khuzistanica* Jamzad essential oil. *Journal of Food Processing and Preservation*, 42 (3), e13557.
- Lai, P.K. & Roy, J. (2004). Antimicrobial and chemo preventive properties of herbs and spices. *Current Medicinal Chemistry*, 11, 1451-1460.
- Mahboubi, M., & Kazempour, N. (2017). Antioxidant and antimicrobial activity of *Satureja khuzistanica* Jamzad essential oil, ethanol and aqueous extracts. *Biharean Biologist*, 1-12.

- Mojaddar-Langroodi A., Tajik H., & Mehdizadeh T. (2019). Antibacterial and antioxidant characteristics of *Zataria multiflora* Boiss essential oil and hydroalcoholic extract of *Rhus coriaria* L. *Journal of Food Quality and Hazards Control*, 6, 16-24.
- Mongelli, E., Pmpuro, S., Coussio, J., Sulomon, H., & Ciccina, G. (2000). Cytotoxic and DNA interaction activities of extracts from medicinal plants used in Argentina. *Journal of Ethnopharmacology*, 71,145-147.
- Mohammadpour, G., Tahmasbpour, R., Noureini, S., Tahmasbpour, E., & Bagherpour, G. (2015). In vitro antimicrobial and cytotoxicity assays of *Satureja bakhtiarica* and *Zataria multiflora* essential oils. *American Journal of Phytomedicine and Clinical Therapeutics*, 3(6), 502-511.
- Naghbi, F., Mosaddegh, M., Mohammadi Motamed, S., & Ghorbani, A. (2005). Labiatae family in folk medicine in Iran: from ethnobotany to pharmacology. *Iranian Journal of Pharmaceutical Research*, 2, 63-79.
- Poon, S. L., McPherson, J. R., Tan, P., Teh, B. T., & Rozen, S. G. (2014). Mutation signatures of carcinogen exposure: genome-wide detection and new opportunities for cancer prevention. *Journal of Genome Medicine* 6:24.
- Raeisi, M., Ghorbani-Bidkorpeh, F., Hashemi, M., Tepe, B., Moghaddam, Z., Aman-Mohammadi, M., & Noori, S.M.A. (2019). Chemical composition and antibacterial and antioxidant properties of essential oils of *Zataria multiflora*, *Artemisia deracunculus* and *Mentha piperita*. *Medical Laboratory Journal*, 13(2), 1-7.
- Raeisi, M., Hashemi, M., Aminzare, M., Afshari, A., Zeinali, T. & Jannat, B. (2018). An investigation of the effect of *Zataria multiflora* Boiss and *Mentha piperita* essential oils to improve the chemical stability of minced meat. *Veterinary world*, 11, 1656-1662.
- Rafieian-Kopaie, M., & Nasri, H. (2015). On the occasion of World Cancer Day 2015: the possibility of cancer prevention or treatment with antioxidants: the ongoing cancer prevention researches. *International Journal of Preventive Medicine*, 6, 108.
- Saeedi-Saravi, S.S., & Shokrzadeh, M. (2008). The study of hepatic and renal disorders in mice which were administered ethyl acetate extract of plant *Sambucus ebulus* Intraperituenally (IP) and effect of vitamins C and E on prevention of its disorders. *Journal of Toxicology Letters*, 180, 57–58.
- Saidi, M. (2014). Antioxidant Activities and Chemical Composition of Essential Oils from *Satureja khuzestanica*, *Oliveria decumbens* and *Thymus daenensis*. *Journal of Essential Oil Bearing Plants*, 17(3), 513-521.
- Sefidkon, F., Naderi, M., Ghushegi, SA. (2013). Comparison of anticancer effects of nanocapsules of *Nasturtium officinalis* (L.) R. Br. extract with methanolic extract and its fractions. *Iranian Journal of Medicinal Aromatic Plants* 29(1): 35-50.
- Stammati, A., Bonsi, P., Zucco, F., Moezelaar, R., Alakomi, H.L. & Von-Wright, A. (1999). Toxicity of selected plant volatiles in microbial and mammalian short-term assays. *Journal of Food Chemistry and Toxicology* 37: 813–823.
- Sylvester, P.W. (2011). Optimization of the tetrazolium dye (MTT) colorimetric assay for cellular growth and viability. *Methods of Molecular Biology*, 716,157-168.
- W.H.O, 2013. WHO traditional medicine strategy: (2014-2023), acces: http://www.who.int/medicines/publications/traditional/trm_strategy14_23/en.
- Yang, H., & Dou, Q.P. (2010). Targeting apoptosis pathway with natural terpenoids: implications for treatment of breast and prostate cancer. *Current Drug Targets*, 11(6), 733-744.
- Zali, H., Rezaei-Tavirani, M. & Azodi, M. (2011). Gastric cancer: prevention, risk factors and treatment. *Gastroenterology Hepatology Bed Bench*, 4(4), 175-185.
- Zamanian-Azodi, M., Rezaie-Tavirani, M., Heydari- Kashal, S., kalantari, S., Dalilan, S. & Zali, H. (2012). Proteomics analysis of MKN45 cell line before and after treatment with Lavender aqueous extract. *Gastroenterology Hepatology Bed Bench*, 5(1), 35-42.
- Zandi-Sohani, N., & Ramezani, L. (2015). Evaluation of five essential oils as botanical acaricides against the strawberry spider mite *Tetranychus turkestanii* Ugarov and Nikolskii. *International Biodeterioration and Biodegradation*, 98, 101-106.
- Zeytinoglu, M., Aydin, S., Ztrk, Y., Hsn, K. & Baser, C. (1998). Inhibitory effects of carvacrol on DMBA induced pulmonary tumorigenesis in rats. *Acta Pharmaceutica Turcica*, 40, 93–98.