

Anatomical and Morphometric Study of Middle Ear Ossicles in Lori Sheep and Native Cattle of Shahrekord Region

Rahmat Allah Fatahian Dehkordi^{1*} , Amirhossein Torabi², Mojgan Sadat Azimi²,
Hanie Mohammadi², Kazem Norozi³

1. Department of Basic Sciences, Faculty of Veterinary Medicine, University of Shahrekord, Shahrekord, Iran.
2. DVM Student, Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, Iran.
3. Expert of Department of Anatomical Sciences, Faculty of Veterinary Medicine, University of Shahrekord, Shahrekord, Iran.

Abstract

BACKGROUND: The ossicles comprise three small bones (malleus, incus, and stapes) known as the smallest bones in the body that play an essential role in transmitting sound to the inner ear.

OBJECTIVES: The study was conducted to compare the structure of the middle ear in native sheep and cattle.

METHODS: To study the details of middle ear ossicles in Lori sheep ($n=5$) and native cattle ($n=5$), bio-anatomical parameters of the malleus, incus, and stapes ossicles were measured. Animals were male and adult. In this study, the ossicles were carefully described by the appropriate devices, and after separation, the ossicles were measured using a caliper and digital scales. Bio-anatomical changes were evaluated under magnification provided by stereomicroscope.

RESULTS: The cow's middle-ear ossicles differed from those of sheep in some details. The results revealed obvious differences in length, width, and thickness of malleus, incus, and stapes in cows compared to sheep. However, further differences were observed in the malleus compared to the other two ossicles when the ossicles of the two species were evaluated. After examining the two studied species, the least difference was observed between the ossicles of the two animals in the stapes.

CONCLUSIONS: To conclude, this study could compare middle ear ossicles between sheep and native cattle in the Shahrekord province. Anyway, it is conceivable that the biometric sizes of the middle ear ossicles may be affected by animal species.

KEYWORDS: Anatomy, Cow, Middle ear, Ossicles, Sheep

Correspondence

Rahmat Allah Fatahian Dehkordi, Department of Basic Sciences, Faculty of Veterinary Medicine, University of Shahrekord, Shahrekord, Iran. Tel: +98 (038) 32324427, Fax: +98 (038) 32324427, Email: fatahian_1349@yahoo.com

Received: 2021-04-06

Accepted: 2021-07-05

Copyright © 2022. This is an open-access article distributed under the terms of the Creative Commons Attribution- 4.0 International License which permits Share, copy and redistribution of the material in any medium or format or adapt, remix, transform, and build upon the material for any purpose, even commercially.

How to Cite This Article

Fatahian Dehkordi, R A ., Torabi, A H., Azimi, M S, Mohammadi H, Norozi K. (2022). Anatomical and Morphometric Study of Middle Ear Ossicles in Lori Sheep and Native Cattle of Shahrekord Region. *Iranian Journal of Veterinary Medicine*, 16(1), 72-80.

Introduction

The performance of hearing is a specific feeling observed among the upper classes of the animal kingdom and is one of the most important sensory inputs to the maintenance of life. It seems very interesting such a small structure like the ear performs a vital act of hearing. There are three small ossicles, called the malleus, incus, and stapes, inside the middle ear (Saha *et al.*, 2017). In the discussion of the middle ear, it has been turned out that it needs to evolve many times; in this way, one ossicle has evolved in primitive birds and reptiles and then in the form of three ossicles in mammals (Tucker, 2017). Wible and Gaudin (2004) reported a lack of processes for muscular fixation in the middle ear in *Euphractus sexcinctus*, but they did not show any degree of muscular development (Wible and Gaudin, 2004). The next experiences were developed from the reptilian condition, which then extended to reptiles and birds and the mammalian middle ear (Pfaff *et al.*, 2019).

The mammalian middle ear is the most basic morphological feature that signalizes this class of vertebrates. Middle ear skeletal pattern differs obviously among vertebrates from those of other amniotes and has attracted the attention of comparative zoologists for years (Meng *et al.*, 2020). By revealing the structural components of the middle ear, examination of the bones involved is not out of reach. The middle ear has three ossicles, the malleus, incus, and stapes (Cañestro *et al.*, 2007; Péus *et al.*, 2020). Researchers have shown more terrestrial mammals can be arranged into two basic middle ear types established upon their ossicular morphologies (also seen in fossils); with a range of intermediate or transitional morphology between both types. These groups have malleus with a long process (processus gracilis) fused to the ectotympanic through the gonial, the wide transverse lamina between the manubrium and the articular surface (for the incus), and the prominent bone mass near the base of the manubrium, which is called the orbicular apophysis (Fleischer, 2013; Wang *et al.*, 2019).

Recent genetic analyses on the mouse have provided information about genes central to middle ear formation (Mallo, 2001, 2003; Fritzsche and Elliott, 2017). However, middle ear bones are essential in mammals for increasing the sensory conduction of

auditory capacity, the way that they conduct sound from the outer ear to the inner ear (Kerber and Sánchez-Villagra, 2019).

Large middle ear ossicles can relatively improve the functional sensitivity of the sensory cells inside the inner ear vestibule to low frequencies with loose ligament attachments and large tympanic membranes (Jurado and Marquardt, 2020). Therefore, one goal from investigations of auditory genetics and physiology is to express the relationship between the middle ear's structure and the function of the auditory signal process. The purposes of the present study were to morphologically describe the middle ear region with particular emphasis on the auditory ossicles; and to carry out a bio-morphometric comparative study of the middle ear ossicles in the two species, *Lori* sheep and *Native* cow.

Materials and Methods

Animals

The study was a biometrical basically sciences-based pragmatic study, performed in the Department of Anatomy in Shahrekord University to monitor the morphological and biometrical differences between ossicles of the middle ear in a period of approximately two months from collecting samples until ossicles separation. A number of cattle and sheep skull bones (in Shahrekord district), each separately and possessing entire ossicular series in the pattern of 'with no discontinuity' and 'without erosion/deterioration' were selected. Bones that seemed to be worn out in consequence of ear disorders and/or complicated diseases were excluded from this investigation. The proposal was approved by the Ethics Committee of Deputy of Faculty and University of Shahrekord approved this research project. The methods used to get different bones of the ear were similar to the early stages of "the method of the mastoidectomy," which was carried out according to a previous protocol (Mitchell and Coulson, 2017). To expose the malleus, incus, and stapes bones, the tympanic bulla was required to be made accessible using a tiny drill. Afterward, the middle ear ossicles were separated from the tympanic cavity and examined using an anatomical stereomicroscope (Model

Olympus Co., Tokyo, Japan). Issues that were considered included; i) description of the anatomical structure of the middle ear ossicles and ii) measurement of the biometric parameters of the middle ear ossicles including weight, thickness, and diameter; comparison of values was made relatively.

Radiographs were established from the middle ear in medial view. The film distance from the X-ray source was about 20 cm and the radiographic device was regulated at 45 kV (KV) and 20 mA/s. An X-ray (Ralco s.r.l Comp) set was used to take a radiograph of the middle ear bones, including malleus, incus, and stapes, on a high-resolution mammography film.

Statistical Analysis

Data of middle ear ossicles were analyzed using the SPSS statistical software package version 23.0.0 for Windows. Data were expressed as mean \pm standard deviation (SD), and statistical variations were tested by the Student-T test. The biometric values of variations, including the weight, length, width, and thickness concerning middle ear ossicles were corrected between two species and the method used was considered with the probability of P -value < 0.05 as statistically significant.

Results

Gross anatomical findings of dissected specimens exhibited that the tympanic bulla included three ossicles of malleus, incus, and stapes (Figures 1 to 3). Figures 3 demonstrate radiographs of middle ear ossicles from lateral and medial views.

The malleus was a hammer-molded small bone that joins to the next bone (incus) and is attached to the inner surface of the tympanic membrane. The incus ossicle, the second bone in the middle ear, was the anvil-shaped small bone that connected to the third middle ear bone (Figures 1 and 2). Stapes, the last bone (Figures 1 and 2), was the stirrup-molded small bone which was placed on other bones, and transported to the oval opening (an opening that leads from the middle ear to the inner ear cavity).

Data from the biometric analysis of three malleus, incus, and stapes bones in both species (sheep and cow) are shown in Table 1. The results revealed an obvious difference in length, width, and thickness of malleus ossicle in the cow compared to the sheep

(Table 1). All the measures except for the weight of malleus ossicle in the cow showed a significant difference ($P < 0.05$). Hence, the weight of malleus ossicle in the cow was not significantly different from that in the sheep ($P > 0.05$). Furthermore, there was a significant difference in terms of biometric dimensions of length, thickness, and width (head, middle, and tail) of the malleus ossicle between both species ($P < 0.05$).




Figure 1. The different parts that make up ossicle; ossicular chain link in sheep mature (dorsal view); M: Handle of malleus; I: Incus; S: Stapes; T: tympanic bulla; M: muscular process; E: external acoustic meatus;

A similar pattern was followed for biometric sizes of the incus ossicle in the middle ear (Table 1), as the biometric dimensions of length and width in the incus ossicle of the cow were significantly higher than those of the sheep ($P < 0.05$). The differences between the two species in terms of weight of incus ossicle did not follow a similar pattern like other biometric values; as there were no significant differences regarding parameters mentioned between two animals, despite the higher weight of incus ossicle in the cow compared to the sheep ($P > 0.05$).




Figure 2. Part x, Ossicles in mature cow; M, Malleus; I, Incus; S, Stapes; A, length of malleus; B, width of malleus; C, thickness Of head in malleus; D, length of incus; E, width of incus; F, length of stapes; G, width of stapes; Part y, Ossicles in mature sheep; I, Incus; M, Malleus; S, Stapes; A, length of malleus; B, width of malleus; C, thickness of head in malleus; D, length of incus; E, width of incus; F, length of stapes; G, width of stapes.




Figure 3. Temporal bone of sheep (1) and cattle (2), tympanic part; radiographs of the middle ear ossicles in sheep (1) and cow (2), medial view. M, malleus; I, incus; S, stapes; T: tympanic bulla; M: muscular process; E: external acoustic meatus.

The values of weight, diameter, and length concerning stapes ossicle in every two animals are presented in [Table 1](#). It should be noted that there was a significant difference between the two species regarding the width of the stapes ossicle ($P < 0.05$). Biometric values of thickness, length, and weight were higher in the cow compared to the sheep during the anatomical evaluation of this bone except for the weight, as there was no significant difference between the cow and sheep in terms of weight of stapes ossicle ($P > 0.05$).

Table 1. Biometric parameter of malleus, incus and stapes between sheep and cow (mm) (Mean \pm SE)

Row	Species	Parameter	Mean \pm SE
1	sheep	malleus.sheep.length	8.97 \pm .363 ^a
2	cow	malleus.cow.length	10.22 \pm .248 ^a
1	sheep	malleus.sheep.head width	2.01 \pm .028 ^b
2	cow	malleus.cow.head width	2.20 \pm .051 ^b
1	sheep	malleus.sheep.tail width	.78 \pm .020 ^c
2	cow	malleus.cow.tail width	.85 \pm .025 ^c
1	sheep	malleus.sheep.middle width	3.49 \pm .024 ^d
2	cow	malleus.cow.middle width	3.95 \pm .015 ^d
1	sheep	malleus.sheep.weight	.029 \pm .001
2	cow	malleus.cow.weight	.040 \pm .004
1	sheep	incus.sheep.length	4.90 \pm .020 ^e
2	cow	incus.cow.length	5.95 \pm .103 ^e
1	sheep	incus.sheep.width	2.42 \pm .017 ^f
2	cow	incus.cow.width	2.83 \pm .049 ^f
1	sheep	incus.sheep.between two branches	3.32 \pm .063
2	cow	incus.cow.between two branches	4.79 \pm .031
1	sheep	incus.sheep.weight	.030 \pm .029
2	cow	incus.cow.weight	.060 \pm .001
1	sheep	stapes.sheep.length	3.44 \pm .065
2	cow	stapes.cow.length	3.62 \pm .005
1	sheep	stapes.sheep.width	2.49 \pm .043 ^g
2	cow	stapes.cow.width	2.69 \pm .014 ^g
1	sheep	stapes.sheep.thickness	.580 \pm .023
2	cow	stapes.cow.thickness	.670 \pm .020
1	sheep	stapes.sheep.weight	.024 \pm .011 ^h
2	cow	stapes.cow.weight	.025 \pm .010 ^h

Significant differences between each two values of a single parameter compared between two species are shown with same letters; otherwise there is no letter denotation.

Discussion

Although some researchers have evaluated the skeletal structure of the middle ear from a descriptive point of view (Erdogan and Kilinc, 2010; Padmini and Rao, 2013; Saha *et al.*, 2017; Simaei *et al.*, 2017), no study has fully compared the middle ear bones (malleus, incus, and stapes) between the two species of cattle and sheep. Therefore, the present study comparatively evaluated these bone structures in the mentioned two species. Findings around biometric values of middle ear ossicles showed a significant difference between the two species ($P < 0.05$). However, no difference was observed regarding the effect of sex on the biometric rate in both animals (data not shown).

The researchers have shown that measuring some of the middle ear ossicles (e.g., stapes) plays an important role in the output impedance of the ear (Grossöhminen *et al.*, 2017). On the other hand, these effects can have another effective function in synthetic hearing aids; in this way, it replaces some lost parts of the body. In another demographic study in India conducted on the morphometric dimensions of male human ear ossicles, Sodhi *et al.* (2017) showed that exact measurements of the ossicles could be very helpful in designing the prosthesis in ossicular chain pathology (Sodhi *et al.*, 2017).

As previously identified, reports revealed no significant difference between middle ear bones in male

and female rabbits. This may lead to the result that diversity in the characteristics of middle ear ossicles, particularly bio-anatomical parameters, does not matter so much in terms of sex (Kurtul *et al.*, 2003), as our results agreed with the results of this study. Although the findings suggest no gender difference in the structure of the middle ear ossicles (Kurtul *et al.*, 2003), in the current study, the effect of gender factor was considered. In some old and new samples of middle ear bones, differences were observed, as Stoessel *et al.* (2016) on the morphology of the middle ear ossicle displayed that there was a significant difference between neandertals and anatomically modern humans (Stoessel *et al.*, 2016).

The gross anatomical structure of the malleus, incus, and stapes ossicles in this study were the same as recent reports (Seibel *et al.*, 2006; Gurr *et al.*, 2011; Kuriakose and Sagar, 2014; Péus *et al.*, 2020). Some studies have shown tangible differences, as Kurtul *et al.* (2003) showed that some middle ear ossicles in rabbits, in anatomical appearance, have an obvious difference with the rest of animals. They found that there was a great variation, especially in the processes and handling in these ossicles (Kurtul *et al.*, 2003).

Regardless of the significant differences between some biometric quantitative values of the middle ear ossicle in this study, the variations indicated a sensible difference in measured values about ossicles of the cattle relative to the sheep. However, with an overview of the "P-value" of the measured number of ossicles, we found that malleus ossicles had the least alteration in both species of cattle and sheep. While in the other two ossicles, namely incus and stapes, the most alteration was observed regarding biometrical measurements. Researches have shown that between the ossicles, the incus is the most fixed ossicle and the stapes ossicle has the most variability as long as their morphological changes are individual and important (Mogra *et al.*, 2014; Noussios *et al.*, 2016; Martonos *et al.*, 2019). Previous studies

have shown that congenital ossicle anomalies are accompanied by facial nerve abnormalities. Furthermore, Padmini and Rao (2013) in evaluating morphological changes of human fetal ear ossicles showed that impoverished human fetuses could be used in homograft forms to substitute corroded mature ear ossicles (Padmini and Rao, 2013).

The tympanic cavity is surrounded by the tympanic membrane, which in most mammalian species forms a glandular, cartilaginous, or bony structure. The structure of the tympanic membrane differs among mammalian species considering the bony main composition or ingredients that contribute to the structure (Ekdale, 2016). Therefore, it seems that the distinct difference between the bones of the two species under this study is closely linked to the dimensions of tympanic bone. In this respect, as is clear, tympanic bone shows a larger dimension in ruminants (Gurr *et al.*, 2011; Péus *et al.*, 2020). Therefore, it is easier to understand the distinct difference between the measured values in the malleus, incus, and stapes ossicles between two animal species; however, some bony values did not show significant differences.

Conclusion

It could be concluded that anatomical parameters of middle ear ossicles examinations in sheep and cow showed some significant differences in ossicles detail, however, these findings can be useful for future studies.

Acknowledgments

This work was financially supported by the University of Shahrekord, Iran. In the end, the authors of the paper, from experts in the field of anatomy of the department of basic sciences thanks and appreciates.

Conflict of Interest

The authors declared no conflict of interest.

References

- Cañestro, C., Yokoi, H., & Postlethwait, J. H. (2007). Evolutionary developmental biology and genomics. *Nature Reviews Genetics*, 8(12), 932-942. [DOI:10.1038/nrg2226] [PMID]
- Ekdale, E. G. (2016). Form and function of the mammalian inner ear. *Journal of Anatomy*, 228(2), 324-337. [DOI:10.1111/joa.12308] [PMID] [PMCID]
- Erdogan, S., & Kilinc, M. (2010). Gross anatomy and arterial vascularization of the tympanic cavity and osseous labyrinth in mid-gestational bovine fetuses. *The Anatomical Record*, 293(12), 2083-2093. [DOI:10.1007/978-3-642-67143-2_2]
- Fleischer, G. (2013). Evolutionary principles of the mammalian middle ear. Springer Science & Business Media. pp, 8-10. [DOI:10.1007/978-3-642-67143-2_2]
- Fritsch, B., & Elliott, K. L. (2017). Evolution and development of the inner ear efferent system: transforming a motor neuron population to connect to the most unusual motor protein via ancient nicotinic receptors. *Frontiers in Cellular Neuroscience*, 11, 114. [DOI:10.3389/fncel.2017.00114] [PMID] [PMCID]
- Grossöhmichen, M., Waldmann, B., Salcher, R., Prenzler, N., Lenarz, T., & Maier, H. (2017). Validation of methods for prediction of clinical output levels of active middle ear implants from measurements in human cadaveric ears. *Scientific Reports*, 7(1), 1-10. [DOI:10.1038/s41598-017-16107-9] [PMID] [PMCID]
- Gurr, A., Pearson, M. D., & Dazert, S. (2011). Lambs' temporal bone anatomy under didactic aspects. *Brazilian Journal of Otorhinolaryngology*, 77(1), 51-57. (Anatomia do osso temporal de ovelhas sob aspectos didáticos.) [DOI:10.1590/S180886942011000100009] [PMID]
- Jurado, C., & Marquardt, T. (2020). On the Effectiveness of airborne infrasound in eliciting vestibular-evoked myogenic responses. *Journal of Low Frequency Noise, Vibration and Active Control*, 39(1), 3-16. [DOI:10.1177/1461348419833868]
- Kerber, L., & Sánchez-Villagra, M. R. (2019). Morphology of the middle ear ossicles in the rodent Perimys (Neoeuplacididae) and a comprehensive anatomical and morphometric study of the phylogenetic transformations of these structures in caviomorphs. *Journal of Mammalian Evolution*, 26(3), 407-422. [DOI:10.1007/s10914-017-9422-9]
- Kuriakose, S., & Sagar, S. (2014). Morphometry and variations of malleus with clinical correlations. *International Journal of Anatomy and Research*, 2(1), 191-194.
- Kurtul, I., Cevik, A., Bozkurt, E. U., & Dursun, N. (2003). A detailed subgross morphometric study on the auditory ossicles of the New Zealand rabbit. *Anatomia, Histologia, Embryologia*, 32(4), 249-252. [DOI:10.1046/j.1439-0264.2003.00483.x] [PMID]
- Mallo, M. (2001). Formation of the middle ear: recent progress on the developmental and molecular mechanisms. *Developmental Biology*, 231(2), 410-419. [DOI:10.1006/dbio.2001.0154] [PMID]
- Mallo, M. (2003). Formation of the outer and middle ear, molecular mechanisms. *Current Topics in Developmental Biology*, 57, 85-113. [DOI:10.1016/S0070-2153(03)57003-X]
- Martonos, C., Damian, A., Gudea, A., Bud, I. T., & Stan, F. (2019). Morphological and morphometrical study of the auditory ossicles in chinchilla. *Anatomia, Histologia, Embryologia*, 48(4), 340-345. [DOI:10.1111/ahel.12446] [PMID]
- Meng, J., Mao, F., Han, G., Zheng, X. T., Wang, X. L., & Wang, Y. (2020). A comparative study on auditory and hyoid bones of Jurassic euharamiyidans and contrasting evidence for mammalian middle ear evolution. *Journal of Anatomy*, 236(1), 50-71. [DOI:10.1111/joa.13083] [PMID]
- Mitchell, S., & Coulson, C. (2017). Endoscopic ear surgery: a hot topic? *The Journal of Laryngology and Otology*, 131(2), 117. [DOI:10.1017/S0022215116009828] [PMID]
- Mogra, K., Gupta, S., Chauhan, S., & Panwar, L. (2014). Morphological and morphometrical variations of malleus in human cadavers. *Journal of Clinical Medicine Research*, 2, 186-192. [DOI:10.5455/2320-6012.ijrms20140562]
- Noussios, G., Chouridis, P., Kostretzis, L., & Natsis, K. (2016). Morphological and Morphometrical Study of the Human Ossicular Chain: A Review of the Literature and a Meta-Analysis of Experience Over 50 Years. *Journal of Clinical Medicine Research*, 8(2), 76-83. [DOI:10.14740/jocmr2369w] [PMID] [PMCID]
- Padmini, M., & Rao, B. (2013). Morphological variations in human fetal ear ossicles-a study. *International Journal of Anatomy and Research*, 1(2), 40-42.

- Péus, D., Dobrev, I., Pfiffner, F., & Sim, J. H. (2020). Comparison of sheep and human middle-ear ossicles: anatomy and inertial properties. *The Journal of Comparative Physiology A*, 206(5), 683-700. [DOI:10.1007/s00359-020-01430-w] [PMID] [PMCID]
- Pfaff, C., Schultz, J. A., & Schellhorn, R. (2019). The vertebrate middle and inner ear: A short overview. *Journal of Morphology*, 280(8), 1098-1105. [DOI:10.1002/jmor.20880] [PMID] [PMCID]
- Saha, R., Srimani, P., Mazumdar, A., & Mazumdar, S. (2017). Morphological Variations of Middle Ear Ossicles and its Clinical Implications. *Journal of Clinical and Diagnostic Research*, 11(1), AC01-AC04. [DOI:10.7860/JCDR/2017/23906.9147]
- Seibel, V. A., Lavinsky, L., & Irion, K. (2006). CT-Scan sheep and human inner ear morphometric comparison. *Brazilian Journal of Otorhinolaryngology*, 72(3), 370-376. [DOI:10.1016/S1808-8694(15)30971-X]
- Simaei, N., Soltananejad, F., Najafi, G., & Jalali, A. S. (2017). Anatomical and morphometrical study of middle ear ossicles in 2 to 3-month-old Makouei sheep fetuses. *Veterinary Research Forum*, 185, 1-11.
- Sodhi, S., Sing, Z., & Lal, J. (2017). Morphometric dimensions of human ear ossicles of males. *National Journal of Medical Research*, 7, 47-51.
- Stoessel, A., David, R., Gunz, P., Schmidt, T., Spoor, F., & Hublin, J. J. (2016). Morphology and function of Neandertal and modern human ear ossicles. *Proceedings of the National Academy of Sciences of the United States of America*, 113(41), 11489-11494. [DOI:10.1073/pnas.1605881113] [PMID] [PMCID]
- Tucker, A. S. (2017). Major evolutionary transitions and innovations: the tympanic middle ear. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372, 1-11. [DOI:10.1098/rstb.2015.0483] [PMID] [PMCID]
- Wang, H., Meng, J., & Wang, Y. (2019). Cretaceous fossil reveals a new pattern in mammalian middle ear evolution. *Nature*, 576(7785), 102-105. [DOI:10.1038/s41586-019-1792-0] [PMID]
- Wible, J. R., & Gaudin, T. (2004). On the cranial osteology of the yellow armadillo *Euphractus sexcinctus* (Dasypodidae, Xenarthra, Placentalia). *Annals of Carnegie Museum*, 73, 117-196.

بررسی تشریحی و ریخت‌سنجی استخوان‌های گوش میانی گوسفند لری و گاو بومی منطقه شهرکرد

رحمت الله فتاحیان دهکردی^{۱*}، امیرحسین ترابی^۲، مژگان سادات عظیمی^۳، هانیه محمدی^۲، کاظم نوروزی^۲

^۱ گروه علوم پایه، دانشکده دامپزشکی، دانشگاه شهرکرد، شهرکرد، ایران

^۲ دانشجوی دوره دکترای عمومی دامپزشکی، دانشکده دامپزشکی، دانشگاه شهرکرد، شهرکرد، ایران

^۳ کارشناس گروه علوم تشریحی، دانشکده دامپزشکی، دانشگاه شهرکرد، شهرکرد، ایران

(دریافت مقاله: ۱۷ فروردین ۱۴۰۰، پذیرش نهایی: ۱۴ تیر ۱۴۰۰)

چکیده

زمینه مطالعه: استخوان‌چه‌های گوش میانی، شامل سه استخوان کوچک بوده و به‌عنوان کوچکترین استخوان‌های بدن شناخته می‌شوند که نقش مهمی در انتقال صدا به گوش داخلی ایفا می‌کنند.

هدف: مطالعه برای مقایسه ساختار گوش میانی در گوسفند و گاو بومی انجام شد.

روش کار: به‌منظور بررسی جزئیات استخوان‌های گوش میانی در گوسفند لری (۵ عدد) و گاو بومی (۵ عدد)، پارامترهای بیو-آناتومیکی استخوان‌چه‌های چکشی، سندان و رکابی اندازه‌گیری شد. حیوانات نر و بالغ بودند. در این مطالعه، استخوان‌چه‌ها با روش‌های دقت توصیف شد و پس از جداسازی آنها، استخوان‌ها با استفاده از کولیس و دستگاه‌های دیجیتال اندازه‌گیری شد. تغییرات بیو-آناتومیکی تحت بزرگ‌نمایی استریومیکروسکوپ بررسی شد.

نتایج: استخوان‌چه‌های گوش میانی گاو با گوسفند در برخی جزئیات متفاوت بود. نتایج نشان داد که تغییرات مشهودی در طول، عرض و ضخامت استخوان‌چه‌های چکشی، سندان و رکابی در گاو نسبت به گوسفند وجود دارد. با این حال، هنگامی که این دو گونه مقایسه شد، تغییرات بیشتری در استخوان چکشی نسبت به دو استخوان دیگر، مشاهده شد. پس از بررسی دو گونه مد نظر، کمترین تغییر قابل توجه بین دو حیوان، در استخوان‌چه رکابی مشاهده شد.

نتیجه‌گیری نهایی: در نتیجه، این مطالعه توانست استخوان‌چه‌های گوش میانی را بین گوسفند و گاو بومی در منطقه شهرکرد با هم مقایسه کند. در هر صورت، قابل تصور است که اندازه‌های بیومتریکی استخوان‌چه‌های گوش میانی ممکن است به‌وسیله‌ی گونه‌های حیوانی تحت تاثیر قرار بگیرد.

واژه‌های کلیدی: استخوان‌چه، آناتومی، گوسفند، گاو، گوش میانی