




## The effect of perturbation training on volleyball players' strength, proprioception and performance

Ebrahim Soltanabadi\*, Maliheh Hadad Nezhad, Mahdi Khaleghi Tazji

Department of Sport Injury and Corrective Exercise, Faculty of Sport Sciences and Health, University of Khazmi, Tehran, Iran. (\*Corresponding author: ✉ [soltanabadiebrahim@yahoo.com](mailto:soltanabadiebrahim@yahoo.com),  <https://www.orcid.org/0000-0002-4401-8347>)

Article Info	Abstract
<p>Original Article</p> <p><b>Article history:</b> Received: 21 May 2022 Revised: 06 August 2022 Accepted: 17 September 2022 Published online: 01 January 2023</p> <p><b>Keywords:</b> functional strength ratio, peak torque, perturbation training, proprioception, shoulder.</p>	<p><b>Background:</b> Perturbation training which is in the context of volleyball routine practice is a type of neuromuscular exercises that is useful in overhead sports; leads to contraction responses through unpredictable perturbation.</p> <p><b>Aim:</b> The present study aims to investigate the effects of perturbation training on volleyball players strength proprioception and performance.</p> <p><b>Materials and Methods:</b> Twenty-four volunteer male volleyball players (age: <math>24.42 \pm 2.15</math> years; height: <math>181.46 \pm 3.00</math>m; Weight: <math>73.87 \pm 3.77</math>kg; BMI: <math>22.43 \pm 0.80</math>) were recruited and randomly assigned into perturbation and control groups, Before the intervention, a set of isokinetic dynamometry was carried out to measure shoulder proprioception as well as rotator cuff muscles strength, what is more, the upper extremity performance was evaluated using Y-Balance Test. For a course of 6 weeks, players of the perturbation group implemented perturbation trainings, 3 times weekly while the control group performed its routine training. The dynamometry was repeated at the end of training period to compare trainings outcome. The Repeated Measures ANOVA statistic was used to find differences between groups.</p> <p><b>Results:</b> A significant difference in shoulder proprioception, as well as the YBT-UQ test, was observed between the groups; Also, the ANOVA showed a significant difference (<math>P &lt; 0.05</math>) in functional strength ratio followed by 6 weeks of perturbation training.</p> <p><b>Conclusion:</b> Perturbation trainings might have benefit to volleyball players conditioning by improvement rotator cuff strength, shoulder proprioception and upper extremity performance. Given this adopting kind of training as a part of regular basic training for volleyball players is highly recommended.</p>

**Cite this article:** Soltanabadi E, Hadad Nezhad M, Khaleghi Tazji M. "The effect of perturbation training on volleyball players' strength, proprioception and performance". *Sport Sciences and Health Research*. 2023, 15(1): 23-33. doi: <https://doi.org/10.22059/SSHR.2023.359173.1090>.



This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY NC), which permits distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.  
EISSN: 2717-2422 | Web site: <https://sshr.ut.ac.ir/> | Email: [sshr@ut.ac.ir](mailto:sshr@ut.ac.ir)

© The Author(s). Publisher: University of Tehran, Faculty of Sport Sciences and Health

## 1. Introduction

Volleyball is a complex and regular sport that requires high technique and tactics and is considered safer than other sports such as football, handball, basketball. The reason for this hypothesis is that in other sports, contact with opposing players is considered part of the game, while volleyball is a non-contact sport [1]. Most volleyball skills such as spike, set, serve, and block require repeated contact of the upper limb with the ball in the overhead position which can cause shoulder malfunction in the long term [2]. Performing repetitive and strong overhead movements will put too much load on the shoulder girdle, causing an imbalance and muscle weakness in the shoulder area [3]. Volleyball is the third most common type of sport including injuries among ball sports [4]. Researchers have also reported shoulder injuries in about 40% of athletic injuries in high school, university, and elite levels [5].

Musculoskeletal injuries are the most common cause of volleyball injuries, with injuries ranging from 1.7 to 10.7 injuries per 1000 hours [6]. Therefore, identifying risk factors and developing strategies for injury prevention are important components of preventive programs in sports injuries [7].

Since volleyball is a non-collision sport, its risk factors may be more related to repetitive overhead movements and changes in biomechanics (i.e., joint mobility, coordination, and disruption of the throwing technique) [8]. Other risk factors include asymmetric tilts and transitional patterns in the shoulder girdle in overhead movements which can cause imbalance and muscle weakness in the shoulder area [3]. Shoulder muscle imbalance is weaker in the external rotator muscles than in the internal rotator. This causes a difference in the range of motion

in the internal and external shoulder joint rotation [9]. Deficiency in proprioception also causes impaired activity control, muscle timing, and joint disorganization [10, 11]. In general, a defect in neuromuscular control is a consequence of injury to the shoulder girdle [12]. Improvement of neuromuscular control and dynamic stability in the shoulder joint requires improvement of the neuromuscular system [13]. Perturbation training can help improve the neuromuscular system because of the variety of exercises and the similarity to the specific needs of the sport [12]. Perturbation training can help facilitate neuromuscular control and speed up the response to unexpected external stimuli [14]. This facilitation of neuromuscular control occurs by facilitating the improvement of partner muscles in motion as well as coordination in the motor chain [15].

Disturbances in overhead movements cause to bring about unpredictable changes in tissue length and muscle contraction response patterns. Perturbation training compared to other overhead exercises helps improve neuromuscular control due to the random nature and the unexpected forces that require both muscle activity preparation and muscle activity response to the stimulus. Performing perturbation exercises at the end of the range of motion improved the efficiency of these exercises because the capsular structures in the range were tightened which increased the sensory signals and increased activity of the muscles involved. In addition, these exercises can improve the sense of articulation [13].

To prevent the problems that a volleyball player may face and the high costs of treating injured people, it is essential to identify those at risk who can be

rescued by providing appropriate preventive training. Muscle strength, proprioception and shoulder stability are the main factors of increasing the athlete's performance. Improving these factors is one of the important goals of coaches and athletes.

Our goal in this study is to investigate the effect of 6 weeks of upper extremity perturbation training on the functional rotator cuff muscle strength, shoulder joint proprioception, and upper body performance in college male volleyball players.

## 2. Materials and Methods

### 2.1. Participation

The present prospective study population consists of academic male volleyball players prone to upper extremity injuries. Out of the whole study population, 24 individuals (Dominant hand: Right; age:  $24.42 \pm 2.15$  years; height:  $181.46 \pm 3.00$ m; weight:  $73.87 \pm 3.77$ kg; BMI:  $22.43 \pm 0.80$ ) were selected as the study sample size according to study inclusion criteria. Participants were then assigned to experimental (perturbation training) and control (no intervention) groups, each including 12 subjects. Inclusion criteria include: 1. Obtain less than 30 points on the CKCUEST test; 2. General health; 3. BMI between 19 and 25; 4.

Be right-handed. Exclusion criteria include: 1. Upper extremity injury in the past 6 months; 2. Missing more than three sessions in practice period; 3. Musculoskeletal abnormalities in the upper extremity; 4. Dissatisfaction with training. The CKCUEST test was used to identify people who were prone to upper limb injury.

### 2.2. Procedure

All subjects completed a brief personal

history form regarding date of birth, number of years of playing, and dominant arm and shoulder injuries. The present study included pre-test and post-test stages. In the pre-test phase, rotator cuff muscle strength and shoulder proprioception were measured by an isokinetic dynamometer. Upper extremity performance was also assessed by the Y Balance Test. Next, the experimental group performed an exercise program for 6 weeks, 3 sessions per week of perturbation training. The control group, on the other hand, only performed their routine pre-season training and did not participate in any regular perturbation training program. After completing the protocol, the post-test started under the same conditions as the pre-test.

#### 2.2.1. Closed Kinetic Chain Upper Extremity Stability Test (CKCUEST) to identify people prone to injury

Two strips with a width of 1.5in were placed parallel to each other 36in apart on the tile floor measured with a standard tape measure. The starting position for the test was one hand on each piece of tape while assuming a push-up position (Figure 1). They had to pass one hand across their body and touch the tape line on the other side (left-hand-right tape cycle and diverse). After touching the hand, it returns to the initial state. Touches were counted as the number of touches by both hands. The total time for each trial was 15 sec. After warming up, the subject performs three tests and rests for 45 sec between each test. Average of the three tests used for data analysis [16].

#### 2.2.2. Evaluation of the rotator Cuff muscles' proprioception using isokinetic system

Determine maximum external rotation (ER) and internal rotation (IR) to assess shoulder joint proprioception. Proprioception is

measured by applying joint-position reproduction, with a target position at 50% ROM. The test started at the initial position (IR or ER) and progressed to the target position, which the participant held for 5 sec to be measured. Variations of  $\pm 5^\circ$  around the target position were allowed. If this variation exceeds, the trial was discarded and repeated. Three tests for each movement direction (ER-IR and IR-ER) perform (6 tests). Just the dominant arm tested. The participants were blindfolded and given task instructions orally by the examiner. The difference between the position reproduced and the position experienced determines individual error [17].

### 2. 2. 3. Assessing isokinetic shoulder rotator cuff muscle strength

The pre-test and post-test of muscular strength with the isokinetic-engaged concentric internal rotator and eccentric external rotator peak torque assessments on the dominant shoulder. According to Wang et al. (2000), an angular velocity of  $120^\circ$  per second was chosen to reduce the risk of injury [18]. For warm-up and familiarization, the subject performed 10 passive contractions. Throughout the assessments, the subjects were seated without their legs touching the ground. A waist strap secures the trunk to the chair. The shoulder is abducted at  $90^\circ$  and the elbow flex at  $90^\circ$  (Figure 2).



**Figure 1.** CKCUEST test to identify people prone to injury



**Figure 2.** Subject positioned for isokinetic strength assessment

The range of motion for the test is set at  $90^\circ$  of external rotation and  $30^\circ$  of internal rotation ( $120^\circ$ ) because exceeding this range increases the risk of injury. First, the internal rotation concentric contraction and then the external rotation eccentric contraction were measured. Subjects each performed five maximal contractions [19].

### 2. 2. 4. Y-balance test

Y balance test performs to evaluate upper

extremity performance. Y Balance Test Kit consists of a stance platform in which three pieces of polyvinyl chloride tube attach in the medial, inferolateral, and superolateral reach directions. The angle between the posterior and anterior pipe is  $135^\circ$ , and between the posterior is  $90^\circ$ . Each pipe mark is in 0.5 cm increments for measurement. To the right-hand reach distance, the subject placed the right hand on a stance plate while the left reached the

medial direction, immediately following the inferolateral and finally in the superolateral. After 3 trials on the right limb, the mean of three attempts was calculated and recorded. For normalization, the reach distance of each direction divides into the length of the upper limb (Distance from the C7 spinous process to the most distal tip of the right middle finger (cm) in 90° shoulder abduction) [20].

$$\text{Total score} = \frac{\text{Medial} + \text{Inferolateral} + \text{Superolateral}}{3 \times \text{Upper limb length}} \times 100$$

### 2. 2. 5. Intervention and Protocol

Perturbation exercises were performed for 6 weeks, 3 sessions per week and 30 min (warmup 5 min, perturbation training 20 min, cool down 5 min) per session. The training protocol included 11 exercises [13] (Figure 3).



Figure 3. Perturbation training

Exercises include forearm supination/pronation, overhead press, waiter’s walk, overhead lunge, windmill, overhead squat, overhead barbell walk, rocker board, wobble board.

### 2. 3. Statistic

Spss (version 22) use to analyze all data. Shapiro-Wilk and 2-factor repeated measures analysis of variance (ANOVA) with one between-subjects factor (group)

and one within-subjects factor (time) were used to examine the normality of the data and inter-group and intra-group changes ( $\alpha \leq 0.005$ ), respectively.

### 3. Results

Shapiro-Wilk test indicated the normality of data in both perturbation and control groups. The anthropometric data showed in Table 1.

Table 1. Characteristics of participants

Variable	(Mean ± SD)	(Mean ± SD)	P-Value
	Perturbation	Control	
Age	24.00 ± 2.13	24.83 ± 2.17	0.353
Weight	75.34 ± 4.25	72.41 ± 2.66	0.055
Height	182.42 ± 3.15	180.50 ± 2.65	0.121
BMI	22.63 ± 0.89	22.23 ± 0.68	0.224
CKCUEST	27.42 ± 1.44	27.33 ± 1.72	0.899

### 3. 1. Isokinetic strength

#### 3. 1. 1. Peak torque values at 120° s<sup>-1</sup>

Mean concentric and eccentric isokinetic peak torque (PT/BW) values for IR and ER showed in Table 2. The results showed that both concentric IR (F[1,22]= 7.017, P= 0.015) and eccentric ER (F[1,22]= 5.268, P= 0.032) peak torque was a significant difference between-subjects in the post-test. There was a significant increase in pre- to post-test (within-subjects) in peak torque values for concentric IR (P<0.0001) and eccentric ER (P<0.0001).

#### 3. 1. 2. Functional strength ratio

Pre-and post-test ratios (eccentric ER strength/ concentric IR strength) showed in Table 2. The ANOVA revealed a statistically significant difference between-subjects for functional strength ratio (F[1,22]= 4.924, P= 0.037) and for within-subject pre- to post-test in the perturbation group (F[1,22]= 69.706; P<0.0001).


#### 3. 1. 3. Proprioception

There was a significant difference between-subjects (F[1,22]= 0.17.495, P<0.0001) and between the pre-and post-test (within-subjects) in the perturbation group (P<0.0001; Figure 4). Results showed that shoulder internal and external proprioception significantly decreased in

the perturbation group.

#### 3. 1. 4. YBT-UQ test

The mean reach distance for all directions of YBT-UQ is present in Table 3. A significant difference was noticed between-subjects in the medial (F[1,22]= 6.159, P=0.021), superolateral (F[1,22]= 5.111, P=0.034), inferolateral (F[1,22]= 4.457, P=0.046), and composite score (F[1,22]= 5.212, P=0.032) and also there was a significant difference within-subjects in medial (F=7.701, P=0.011), superolateral (F=51.229, P<0.0001), inferolateral (F=344.049, P<0.0001) and composite score (F=166.764, P<0.0001) of YBT-UQ test.



**Figure 4.** Pre- to post-test mean joint position sense (degree) between the 2 groups. PRO= proprioception; pre = pre-test; post = post-test. \* Significant difference (P< 0.05) from PRO pre-test

**Table 2.** Mean concentric and eccentric peak torque values (N.M/BW) for internal rotation (IR) and external rotation (ER) of the dominant shoulder at 120°.s<sup>-1</sup>. CON = concentric IR; ECC = eccentric ER.

	Mean ± SD		F	Sig. (2-tailed)
	Pre test	Post Test		
Concentric IR pre-post test	52.808 ± 2.656	53.883 ± 3.159	240.385	0.000
Eccentric ER pre-post test	36.579 ± 1.666	36.967 ± 1.713	17.807	0.000

\* Significant difference (P< 0.05) from CON IR pre-test value. † Significant difference (P< 0.05) from ECC ER pre-test value

**Table 3.** Functional (external rotation/ internal rotation) strength ratios at 120°.s<sup>-1</sup> in N.m\*

	Pre EX/IR ratio 120°.s <sup>-1</sup>	Post EX/IR ratio 120°.s <sup>-1</sup>
Perturbation	0.693 ± 0.010	0.678 ± 0.009
Control	0.693 ± 0.009	0.695 ± 0.008

\* Pre = pre-test; post = post-test; ER = External Rotation; IR = Internal Rotation

**Table 4.** Mean reach distance of YBT-UQ in Cm\*

Group		Pre	Post
Perturbation	M	103.99 ± 1.89	104.88 ± 1.03
	Sly	74.88 ± 2.29	77.62 ± 1.94
	IL	83.53 ± 3.62	88.49 ± 3.20
	C	93.55 ± 1.23	96.63 ± 1.47
Control	M	103.09 ± 1.30	103.06 ± 1.23
	SL	74.53 ± 1.71	74.32 ± 2.17
	IL	83.67 ± 1.75	83.69 ± 1.80
	C	93.84 ± 1.48	93.75 ± 1.50

\* Pre = pre-test; post = post-test; M = Medial direction; SL = Superolateral; IL = Inferolateral; C= Composite Score

#### 4. Discussion

The present study aim is to investigate the effect of 6 weeks of upper extremity perturbation training on the functional rotator cuff muscle strength, shoulder joint proprioception, and upper body performance in college male volleyball players. Twenty-four volleyball players are prone to upper limb injuries were studied. The results showed that the concentric strength of the internal rotators and the eccentric strength of the external rotators increased, and the functional strength ratio of the rotator cuff muscles decreased after 6 weeks in the experimental group.

Most researches have focused on improving muscle strength, which is known as the basis of physical activity. Exercises such as weight-based exercises [21], ballistic and explosive resistance exercises [22], electrical stimulation exercises [23], and vibration exercises [24] have been used to improve and increase strength. Reasons to improve strength include the adaptation of the neuromuscular system such as increased firing frequency, improved coordination of motor units, increased excitability of motor units, and increased motor unit flow [25]. The shoulder is a joint with a high range of motion. To generate force in the joint needs sufficient stability. External and internal rotator muscles are the most important muscles of the shoulder area

to stabilize shoulder joint movements. To keep the humerus head in the glenoid cavity, a balance of a couple of forces between the external and internal rotators is required; hence, in assessing upper extremity strength, measuring the ratio of external and internal rotator power is the most important factor for injury prevention. Some researchers have reported a normal range of FDR ranging from 0.66 to 1.54 and a BR range between 0.46 and 1.05, which varies depending on the type of test. FDR is the ratio of the eccentric strength of the external rotator muscles to the concentric contraction strength of the internal rotator muscles, whereas BR is the ratio of the eccentric contraction strength of the rotator cuff muscles. Ellenbecker et al. (2004) reported a normal range of shoulder functional strength ratio from 0.6 to 0.75 [26]. If this ratio is less than 0.6 or greater than 0.75, it is considered abnormal. In this case, the muscles become imbalanced, and the risk of shoulder injury increases. Perturbation during overhead activity can cause changes in movement patterns through unexpected changes in muscle length and quick response to muscle contraction [27]. During training, joint forces balance due to improvements in a couple of forces [28]. Improvement in joint forces increases the efficiency of neuromuscular control and better dynamic

stability than the dependence on non-contractile tissues involved in static stability [27, 28]. Perturbation exercises are more effective in neuromuscular control than other overhead exercises because of randomness and unpredictable movements [15]. Therefore, these exercises can be useful for improving movement patterns, enhancing function, and injury prevention.

After 6 weeks, perturbation training showed a significant effect on the proprioception of the shoulder joint in the experimental group. The findings indicated a decrease and improvement in shoulder joint proprioception, which is consistent with previous studies [17, 29, 30, 31].

Swanik et al. (2002) found that plyometric exercises improved shoulder sensation. The peripheral and central nervous adaptations that occur with plyometric exercises improve joint proprioception. These adaptations may be due to repeated stimulation of the mechanical receptors around the joint at the end of the range of motion during exercise. Quick length-tension changes of tendon-muscle structures during eccentric loads facilitate the spindle and Golgi's tendon adaptation. Muscle stretching increases the sensitivity of the muscle spindle, which increases the cooperation of sensory nerves with the central nervous system and finally improves depression [31].

Proprioception receptors identify changes in joint angle, muscle length, and muscle strength. The muscle spindle is one of the proprioceptive receptors. Muscle spindles regulate the activity of alpha motor neurons and participate in tensile reflexes by responding to changes in muscle length or tissue elongation ratio. In this study, the limb position was selected for training at the end of the range of motion because it would challenge proprioception and motor control.

Janwantanakul et al. (2001) reported that joint proprioception improved at the end of the ROM more than at other positions. At the end of the range of motion, the joint capsule is stretched, and the input sensory impulses increase, which activates the muscles further [11].

The upper extremity function evaluates by the Y-balance test. The results showed a significant increase in distance reach in all directions of the Y-balance test after 6 weeks of perturbation training in the experimental group. Most studies have compared the upper extremity function of athletes in different sports or injuries. Upper extremity performance can evaluate by various tests such as Y-balance, CKCUEST, medicine ball throw, etc. Research evaluating upper-extremity performance with Y-balance was scarce.

Fathi et al. (2019) found that strength and plyometric training improved throw performance in adult volleyball players. They suggested strength or plyometric exercises to volleyball-specific exercises to improve performance [32].

In another study, Ignjatovic et al. (2012) showed that resistance training improved the distance of throwing the medicine ball and increased performance [33]. Shoulder movements are the result of a complex process of synthesizing central and peripheral nervous system information for the design, coordination, and synergy of afferent and efferent neurons in feedback and feedforward processes [34]; hence, the performance of overhead athletes requires a static and dynamic stabilization mechanism for coordinating the glenohumeral and thoracic joints to produce an appropriate motor response [11]. Perturbation training facilitates neuromuscular control increases the speed of response to disturbances, and can increase upper extremity function [14].



Facilitation of neuromuscular control probably occurs through the facilitation of the recovery of partner muscles in motion as well as coordination in the kinetic chain from the proximal to the distal limbs [15]. Perturbation may improve coordination in muscle contraction, promote function, and reduce injury [27].

## 5. Conclusions

Due to the effect of perturbation training on performance, strength, and proprioception, these exercises recommend to coaches and athletes whose upper extremity performance is impaired or at risk of injury.

## Conflict of interest

The authors declared no conflicts of interest.

## Authors' contributions

All authors contributed to the original idea, study design: Writing, Review and Editing [all authors], Methodology [all authors], Supervision [Maliheh Hadad Nezhad], Advisor [Mahdi Khaleghi Tazji].

## Ethical considerations

The authors have completely considered ethical issues, including informed consent, plagiarism, data fabrication, misconduct, and/or falsification, double publication and/or redundancy, submission, etc.

Ethical Code: (IR-KHU.KRC.1000.129)

## Data availability

The dataset generated and analyzed during the current study is available from the corresponding author on reasonable request.

## Funding

The article is extracted from the thesis supported by Kharazmi University.

## Acknowledgment

I would like to thank my supervisor and advisor for helping me in this research.

## References

- [1] Bere T, Kruczynski J, Veintimilla N, Hamu Y, Bahr R. "Injury risk is low among world-class volleyball players: 4-year data from the FIVB Injury Surveillance System". *British Journal of Sports Medicine*. 2015; 49(17): 1132-7. <http://dx.doi.org/10.1136/bjsports-2015-094959>.
- [2] Reeser JC, Joy EA, Porucznik CA, Berg RL, Colliver EB, Willick SE. "Risk factors for volleyball-related shoulder pain and dysfunction". *Pm&r*. 2010; 2(1): 27-36. <https://doi.org/10.1016/j.pmrj.2009.11.010>.
- [3] Grabara M. "Comparison of posture among adolescent male volleyball players and non-athletes". *Biology of Sport*. 2015; 32(1): 79-85. <https://dx.doi.org/10.5604%2F20831862.1127286>.
- [4] Seminati E, Minetti AE. "Overuse in volleyball training/practice: A review on shoulder and spine-related injuries". *European Journal of Sport Science*. 2013; 13(6): 732-43. <https://doi.org/10.1080/17461391.2013.773090>.
- [5] Dick R, Sauers EL, Agel J, Keuter G, Marshall SW, McCarty K, et al. "Descriptive epidemiology of collegiate men's baseball injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2003–2004". *Journal of Athletic Training*. 2007; 42(2): 183. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1941283/>.
- [6] Cools AM, Reeser JC. "Shoulder injuries in volleyball". *Handbook of Sports Medicine and Science: Volleyball*. 2017: 93-108.
- [7] Finch C. "A new framework for research leading to sports injury prevention". *Journal of Science and Medicine in Sport*. 2006; 9(1-2): 3-9. <https://doi.org/10.1016/j.jsams.2006.02.009>.
- [8] Vincent HK, Zdziarski LA, Vincent KR. "Review of lacrosse-related musculoskeletal injuries in high school and collegiate players". *Sports Health*. 2015; 7(5): 448-51. <https://doi.org/10.1177/1941738114552990>.
- [9] James LP, Kelly VG, Beckman EM. "Injury risk management plan for volleyball athletes". *Sports Medicine*. 2014; 44: 1185-95. <https://doi.org/10.1007/s40279-014-0203-9>.
- [10] Carpenter JE, Blasler RB, Pellizzon GG. "The

- effects of muscle fatigue on shoulder joint position sense". *The American Journal of Sports Medicine*. 1998; 26(2): 262-5. <https://doi.org/10.1177/03635465980260021701>.
- [11] Janwantanakul P, Magarey ME, Jones MA, Dansie BR. "Variation in shoulder position sense at mid and extreme range of motion". *Archives of Physical Medicine and Rehabilitation*. 2001; 82(6): 840-4. <https://doi.org/10.1053/apmr.2001.21865>.
- [12] Saeterbakken AH, Andersen V, Behm DG, Krohn-Hansen EK, Smaamo M, Fimland MS. "Resistance-training exercises with different stability requirements: time course of task specificity". *European Journal of Applied Physiology*. 2016; 116: 2247-56. <https://doi.org/10.1007/s00421-016-3470-3>.
- [13] Burne G, Tack C. "Overhead perturbation training". *Strength & Conditioning Journal*. 2017; 39(3): 87-93. [doi: 10.1519/SSC.0000000000000280](https://doi.org/10.1519/SSC.0000000000000280).
- [14] Wilk KE, Arrigo CA, Hooks TR, Andrews JR. "Rehabilitation of the overhead throwing athlete: there is more to it than just external rotation/internal rotation strengthening". *Pm&r*. 2016; 8(3): S78-S90. <https://doi.org/10.1016/j.pmrj.2015.12.005>.
- [15] Myers JB, Lephart SM. "The role of the sensorimotor system in the athletic shoulder". *Journal of Athletic Training*. 2000; 35(3): 351. <http://www.ncbi.nlm.nih.gov/pmc/articles/pmc1323397/>.
- [16] Lagniaux F, Juin G, Rambaud A, Thoreux P. "Reference values for the closed kinetic chain upper extremity stability test for elite handball players". *International Journal of Sports Physical Therapy*. 2019; 14(6). <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2953295/>.
- [17] Salles JI, Velasques B, Cossich V, Nicoliche E, Ribeiro P, Amaral MV, et al. "Strength training and shoulder proprioception". *Journal of Athletic Training*. 2015; 50(3): 277-80. <https://doi.org/10.4085/1062-6050-49.3.84>.
- [18] Wang H-K, Macfarlane A, Cochrane T. "Isokinetic performance and shoulder mobility in elite volleyball athletes from the United Kingdom". *British Journal of Sports Medicine*. 2000; 34(1): 39-43. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1724158/>.
- [19] Niederbracht Y, Shim AL, Sloniger MA, Paternostro-Bayles M, Short TH. "Effects of a shoulder injury prevention strength training program on eccentric external rotator muscle strength and glenohumeral joint imbalance in female overhead activity athletes". *The Journal of Strength & Conditioning Research*. 2008; 22(1): 140-5. <https://doi.org/10.1519/jsc.0b013e31815f5634>.
- [20] Gorman PP, Butler RJ, Plisky PJ, Kiesel KB. "Upper quarter Y balance test: Reliability and performance comparison between genders in active adults". *The Journal of Strength & Conditioning Research*. 2012; 26(11): 3043-8. <https://doi.org/10.1519/jsc.0b013e3182472fdb>.
- [21] Wilson GJ, Murphy AJ, Giorgi A. "Weight and plyometric training: Effects on eccentric and concentric force production". *Canadian Journal of Applied Physiology*. 1996; 21(4): 301-15. <https://doi.org/10.1139/h96-026>.
- [22] Newton RU, Rogers RA, Volek JS, Häkkinen K, Kraemer WJ. "Four weeks of optimal load ballistic resistance training at the end of season attenuates declining jump performance of women volleyball players". *The Journal of Strength & Conditioning Research*. 2006; 20(4): 955-61. <https://doi.org/10.1519/r-5050502x.1>.
- [23] Malatesta D, Cattaneo F, Dugnani S, Maffiuletti NA. "Effects of electromyostimulation training and volleyball practice on jumping ability". *The Journal of Strength & Conditioning Research*. 2003; 17(3): 573-9. [https://doi.org/10.1519/1533-4287\(2003\)017%3C0573:eoetav%3E2.0.co;2](https://doi.org/10.1519/1533-4287(2003)017%3C0573:eoetav%3E2.0.co;2).
- [24] Cardinale M, Lim J. "Electromyography activity of vastus lateralis muscle during whole-body vibrations of different frequencies". *The Journal of Strength & Conditioning Research*. 2003; 17(3): 621-4. PMID: 12930196.
- [25] De Villarreal ES-S, Requena B, Newton RU. "Does plyometric training improve strength performance? A meta-analysis". *Journal of Science and Medicine in Sport*. 2010; 13(5): 513-22. <https://doi.org/10.1016/j.jsams.2009.08.005>.
- [26] Davies GJ, Ellenbecker TS, Andrews J, Harrelson G, Wilk K. "Application of isokinetics in testing and rehabilitation". *Physical Rehabilitation of the Injured Athlete*. 2004; 4. <https://www.ncbi.nlm.nih.gov/pubmed/16558647>.
- [27] Guido Jr JA, Stemm J. "Reactive neuromuscular training: A multi-level approach to rehabilitation of the unstable shoulder". *North American Journal of Sports Physical Therapy*. 2007; 2(2): 97.

- <https://www.ncbi.nlm.nih.gov/pubmed/21522207>.
- [28] Cluff T, Crevecoeur F, Scott SH. "A perspective on multisensory integration and rapid perturbation responses". *Vision Research*. 2015; 110: 215-22. <https://doi.org/10.1016/j.visres.2014.06.011>.
- [29] Başkurt Z, Başkurt F, Gelecek N, Özkan MH. "The effectiveness of scapular stabilization exercise in the patients with subacromial impingement syndrome". *Journal of Back and Musculoskeletal Rehabilitation*. 2011; 24(3): 173-9. <https://pubmed.ncbi.nlm.nih.gov/21849731/>.
- [30] Salles JI, Guimarães J, Filho G, Morrissey D. "Effect of a specific exercise strategy on strength and proprioception in volleyball players with infraspinatus muscle atrophy". *Scandinavian Journal of Medicine & Science in Sports*. 2018; 28(9): 2093-9. <https://pubmed.ncbi.nlm.nih.gov/29772095/#article-details>.
- [31] Swanik KA, Lephart SM, Swanik CB, Lephart SP, Stone DA, Fu FH. "The effects of shoulder plyometric training on proprioception and selected muscle performance characteristics". *Journal of Shoulder and Elbow Surgery*. 2002; 11(6): 579-86. <https://pubmed.ncbi.nlm.nih.gov/12469083/>.
- [32] Fathi A, Hammami R, Moran J, Borji R, Sahli S, Rebai H. "Effect of a 16-week combined strength and plyometric training program followed by a detraining period on athletic performance in pubertal volleyball players". *The Journal of Strength & Conditioning Research*. 2019; 33(8): 2117-27. <https://pubmed.ncbi.nlm.nih.gov/29401199/>.
- [33] Ignjatovic AM, Markovic ZM, Radovanovic DS. "Effects of 12-week medicine ball training on muscle strength and power in young female handball players". *The Journal of Strength & Conditioning Research*. 2012; 26(8): 2166-73. <https://pubmed.ncbi.nlm.nih.gov/22027860/>.
- [34] Diedrichsen J, Shadmehr R, Ivry RB. "The coordination of movement: Optimal feedback control and beyond". *Trends in Cognitive Sciences*. 2010; 14(1): 31-9. <https://pubmed.ncbi.nlm.nih.gov/20005767/>.